


Nonequilibrium version of the Einstein relation


Daniel Hurowitz, Ben-Gurion University






D. Hurowitz and D. Cohen, Phys. Rev. E 90, 032129 (2014)

Overview: Nonequilibrium steady state of **sparse** systems


Energy absorption



Nontrivial topology



Sinai regime - current changes sign [2]



Current distribution reflects barrier statistics [3]

Quantum saturation of effective temperature [1]

[1] D. Hurowitz and D. Cohen, *Europhysics Letters*, **93**, 60002 (2011)

[2] D. Hurowitz, S. Rahav and D. Cohen, *Europhysics Letters*, **98**, 20002 (2012)

[3] D. Hurowitz, S. Rahav and D. Cohen, *Phys. Rev. E*, **88**, 062141 (2013)

Brownian motion

The Einstein-Smoluchowski Relation (ESR):

$$D = \mu k_B T, \quad k_B = 1$$

Relation between mobility (μ) and diffusion (D) reflecting microscopics (k_B) in universal way.

This is a special case of a **fluctuation-dissipation relation** between first and second moments.

Drift: $\langle x \rangle = vt, \quad v = \mu F$

Diffusion: $\text{Var}(x) = 2Dt$

ESR: $\frac{v}{D} = \frac{F}{T} \equiv s = \text{affinity (linear response)}$

$s \equiv$ entropy-production-per-distance

FDT is valid close to equilibrium.

To what extent does the ESR hold?

Can it be derived from the NFT?

Non-equilibrium version?

Sinai spreading

Stochastic field: $\mathcal{S}_n \equiv \ln \left[\frac{\vec{w}_n}{\overleftarrow{w}_n} \right]$, $\sigma = \sqrt{\text{Var}(\mathcal{S}_n)}$

Stochastic Motive Force: $\mathcal{S}_\circ = \sum_{n \in \text{ring}} \ln \left[\frac{\vec{w}_n}{\overleftarrow{w}_n} \right]$

If $\frac{\vec{w}_n}{\overleftarrow{w}_n} = \exp \left[-\frac{E_n - E_{n-1}}{T} \right] \rightsquigarrow \mathcal{S}_\circ = 0$

Affinity: $s = \frac{\mathcal{S}_\circ}{N}$

For small s [1]:

Sub-diffusive spreading $x \sim [\log(t)]^2$,



Exponentially small drift $v \sim e^{-\sqrt{N}}$.

For arbitrary s [2,3]:

Complicated expressions for v and D .

For a periodic lattice, no disorder:

$$\frac{v}{D} = \frac{2}{a} \tanh \left(\frac{as}{2} \right)$$





[1] Sinai (1982)

[2] Derrida (1983)

[3] Aslangul, Pottier, Saint-James (1989)

ESR is violated for large s

Observations for finite N



Generalized ESR for a given disorder σ

$$\frac{v}{D} = \frac{2}{a_s} \tanh \frac{a_s s}{2}$$

- (1) For small values of s we have $v/D = s$, in consistency with the ESR.
- (2) For no disorder ($\sigma = 0$) we have $a_s = 1$, reflecting the discreteness of the lattice.
- (3) For finite disorder and moderate s we have $a_s \sim N$, reflecting the length of the unit cell.
- (4) For finite disorder and large s we have $a_s = a_\infty$, reflecting the disorder σ .
- (5) As N becomes larger our results approach those of [2,3], which we call "Sinai step".

Outline

General s dependence

$$\frac{v}{D} = \frac{2}{a_s} \tanh \frac{a_s s}{2}$$

$a_s : N \dots a_\infty$

Poisson ($s \rightarrow \infty$)

$$\frac{v}{D} = \frac{2}{a_\infty}$$

$a_\infty(\sigma) : 1 \dots N$

ESR ($s \rightarrow 0$)

$$\frac{v}{D} = s$$

Given $(1, N, \sigma)$

$a_s = ?$

σ is the log-width of the stochastic field distribution

Nonequilibrium Fluctuation Theorem (NFT) derivation of the ESR

Define x as the winding number times the length of the ring.

$$\frac{P[x(-t)]}{P[x(t)]} = \exp[-\mathcal{S}[x]] \quad \rightsquigarrow \quad \frac{p(-x; t)}{p(x; t)} = e^{-sx}$$

Gaussian approximation (Central Limit Theorem)

$$p(x; t) \approx \bar{p}(x; t) = \frac{1}{\sqrt{4\pi Dt}} \exp\left[-\frac{(x - vt)^2}{4Dt}\right] \quad \rightsquigarrow \quad \frac{v}{D} = s$$

Does the ESR really hold?

NFT and coarse graining

Asymmetric random walk traversing a distance $x = X_1 + \dots + X_{\mathcal{N}}$

$$P(X = +1) = p \equiv \vec{w}\tau$$

$$P(X = -1) = q \equiv \overleftarrow{w}\tau$$

$$P(X = 0) = 1 - p - q$$

Moment generating function $Z(k) = \langle e^{-ikx} \rangle = \left[pe^{-ik} + qe^{+ik} + (1 - p - q) \right]^{\mathcal{N}}$

In the continuous time limit $p, q \ll 1$, $\ln Z(k) = \mathcal{N} \left[pe^{-ik} + qe^{+ik} - (p + q) \right] + \mathcal{O}(\mathcal{N}\tau^2)$

Accordingly, one obtains:

$$p(x; t) = \int_{-\infty}^{\infty} dk e^{ikx + (\vec{w}e^{-ik} + \overleftarrow{w}e^{ik} - (\vec{w} + \overleftarrow{w}))t} \quad \text{satisfies NFT}$$

Correct application of the CLT:

$$\bar{p}(x; t) = \int_{-\infty}^{\infty} dk e^{ik(x - (\vec{w} - \overleftarrow{w})t) - \frac{k^2}{2}(\vec{w} + \overleftarrow{w})t + \cancel{\mathcal{O}(k^3 t)}} = \frac{1}{\sqrt{4\pi Dt}} \exp \left[-\frac{(x - vt)^2}{4Dt} \right]$$

$$v = \vec{w} - \overleftarrow{w}, \quad D = \frac{1}{2}(\vec{w} + \overleftarrow{w}) \quad \rightsquigarrow \quad \frac{v}{D} = \bar{s} = \frac{2}{a} \tanh \frac{as}{2} \quad \text{The affinity is renormalized!}$$

The naive reasoning, based on CLT, is **wrong**, If we smear $p(x)$ we get

$$\frac{\bar{p}(-x; t)}{\bar{p}(x; t)} = e^{-\bar{s}x}$$

Recipe for computing v and D on a periodic array

Dynamics determined by rate equation: $(d/dt)\mathbf{p} = W\mathbf{p}$

W is not symmetric yet periodic, thus Bloch's theorem applies.

Reduced equation for the eigenmodes $\mathbf{W}(\varphi)\psi = -\lambda\psi$, where $\mathbf{W}(\varphi)$ is an $N \times N$ matrix.

Bloch's theorem: $\psi_{n+N} = e^{i\varphi}\psi_n$, where n is the site index mod(N).

Bloch quasi-momentum $\varphi \equiv kN$.

Diagonalizing $\mathbf{W}(\varphi) \rightsquigarrow \{|k, \nu\rangle, -\lambda_\nu(k)\}$, where ν is the band index.

Time dependent solution of the rate equation

$$p_n(t) \approx \frac{1}{L} \sum_{k, \nu} C_{k, \nu} e^{-\lambda_\nu(k)t} e^{ikn} \quad \text{where } C_{k, \nu} \text{ depend on initial conditions.}$$

In the long time limit only λ_0 survives

$$v = i \left. \frac{\partial \lambda_0(k)}{\partial k} \right|_{k=0}$$
$$D = \frac{1}{2} \left. \frac{\partial^2 \lambda_0(k)}{\partial k^2} \right|_{k=0}$$

The Poisson Limit ($s \rightarrow \infty$)

The limit $s \rightarrow \infty$ corresponds to a uni-directional random walk traversing a distance $x = X_1 + \dots + X_N$

$$P(X_n = 1) = w_n \tau$$

$$P(X_n = 0) = 1 - w_n \tau$$

$$P(X_n = -1) = 0$$

Characteristic polynomial for eigenvalues of $\mathbf{W}(\varphi)$


$$\det(\lambda + \mathbf{W}(\varphi)) = \prod_{n=1}^N (\lambda - w_n) + e^{-i\varphi} \prod_{n=1}^N w_n = 0$$

Expanding to second order in λ and φ

$$\lambda = -i \left[\left(\sum_{n=1}^N \frac{1}{w_n} \right)^{-1} \right] \varphi + \frac{1}{2} \left[\left(\sum_{n=1}^N \frac{1}{w_n} \right)^{-3} \left(\sum_{n=1}^N \frac{1}{w_n^2} \right) \right] \varphi^2 + \mathcal{O}(\varphi^3)$$



From the recipe for v and D :

$$a_\infty = \left(\frac{2D}{v} \right)_{s \rightarrow \infty} = \left[\frac{\langle (1/\vec{w})^2 \rangle}{\langle (1/\vec{w}) \rangle^2} \right] = [\text{For log-box distribution}] = \frac{\sigma}{2} \coth \left(\frac{\sigma}{2} \right)$$



Effective lattice constant ($N = 6$)

Reminder: Observations for finite N



Generalized ESR for a given disorder σ

$$\frac{v}{D} = \frac{2}{a_s} \tanh \frac{a_s s}{2}$$

- (1) For small values of s we have $v/D = s$, in consistency with the ESR.
- (2) For no disorder ($\sigma = 0$) we have $a_s = 1$, reflecting the discreteness of the lattice.
- (3) For finite disorder and moderate s we have $a_s \sim N$, reflecting the length of the unit cell.
- (4) For finite disorder and large s we have $a_s = a_\infty$, reflecting the disorder σ .
- (5) As N becomes larger our results approach those of [2,3], which we call "Sinai step".

Spreading analysis and the "Sinai step"

$$\left\langle \left(\frac{\overleftarrow{w}}{\overrightarrow{w}} \right)^\mu \right\rangle \equiv e^{-(s-s_\mu)\mu} \quad [\text{defines } s_\mu]$$

The values $s_{1/2}$, s_1 and s_2 determine crossover points between transport regimes.

For $s = 0$, anomalous time dependent spreading [Sinai],

$$x \sim [\log(t)]^2 \quad \rightsquigarrow \quad v \sim e^{-\sqrt{N}}$$

For finite $s < s_1$ [Bouchaud, Comtet, Georges, Le Doussal, 1987],

$$x \sim t^\mu \quad [\mu \text{ is the value for which } s_\mu = s]$$

Time required to drift $x \sim N$ is $t \sim N^{1/\mu}$, hence

$$v \sim \frac{x}{t} \sim \left(\frac{1}{N} \right)^{\frac{1}{\mu}-1}$$

Crossover at $s = s_{1/2}$ from sub-Ohmic to super-Ohmic behaviour .

For large $s > s_1$ and $N \rightarrow \infty$ [Derrida],

$$v_s = \frac{1 - \langle (\overleftarrow{w}/\overrightarrow{w}) \rangle}{\langle (1/\overrightarrow{w}) \rangle} = \left[1 - e^{-(s-s_1)} \right] v_\infty$$

The affinity dependent length scale a_s


From "Derrida" we have an expression for v in the $N \rightarrow \infty$ limit.

From our reasoning we have in general

$$\frac{v}{D} = \frac{2}{a_s} \tanh \frac{a_s s}{2} \quad \text{with some } a_s.$$

By "reverse engineering" we deduce

$$\begin{cases} a_s \sim N, & s < s_2 \\ a_s \approx \frac{a_\infty}{1 - \langle (\frac{\overleftarrow{w}}{\overrightarrow{w}})^2 \rangle} = \frac{a_\infty}{1 - e^{-2(s-s_2)}}, & s > s_2 \end{cases}$$



s regime	$[0, 1/N]$	$[1/N, s_{1/2}]$	$[s_{1/2}, s_1]$	$[s_1, s_2]$	$[s_2, \infty]$
a_s	irrelevant	$a_s \sim N$			$a_s \approx [1 - e^{-2(s-s_2)}]^{-1} a_\infty$
v_s	$v = 2D s$	$\sim (\frac{1}{N})^{\frac{1}{\mu}-1}$		$v_s \approx [1 - e^{-(s-s_1)}] v_\infty$	
D	$\sim \exp(-\sqrt{N})$	$\sim (\frac{1}{N})^{\frac{1}{\mu}-2}$	$\sim (N)^{2-\frac{1}{\mu}}$	$\sim N$	$D = \frac{1}{2} a_s v_s$

Summary

To what extent does the ESR hold?

As long as $s < 1/N$.

Can it be derived from the NFT?

Yes, provided s is replaced by coarse grained \bar{s} .

Non-equilibrium version?

$$\left\{ \begin{array}{ll} v \sim \left(\frac{1}{N}\right)^{\frac{1}{\mu}-1}, & s < s_1 \\ v \approx [1 - e^{-(s-s_1)}] v_\infty & s > s_1 \end{array} \right.$$

$$\left\{ \begin{array}{ll} a_s \sim N, & s < s_2 \\ a_s \approx \frac{a_\infty}{1 - \langle (\frac{\overleftarrow{w}}{\overrightarrow{w}})^2 \rangle} = \frac{a_\infty}{1 - e^{-2(s-s_2)}}, & s > s_2 \end{array} \right.$$

$$\frac{v}{D} = \frac{2}{a_s} \tanh \frac{a_s s}{2}$$

Epilog: Experiments

- S.H. Lee and D. G. Grier, **Giant Colloidal Diffusivity on Corrugated Optical Vortices**, PRL 2006
- Giorgio Volpe, Giovanni Volpe, Sylvain Gigan, **Brownian Motion in a Speckle Light Field: Tunable Anomalous Diffusion and Selective Optical Manipulation**, Scientific Reports 2014
- S. Nakamura, Y. Yamauchi, M. Hashisaka, K. Chida, K. Kobayashi, T. Ono, R. Leturcq, K. Ensslin, K. Saito, Y. Utsumi, A.C. Gossard, **Nonequilibrium fluctuation relations in a quantum coherent conductor**, PRL 2010
- B. Kung, C. Rossler, M. Beck, M. Marthaler, D. S. Golubev, Y. Utsumi, T. Ihn, and K. Ensslin **Irreversibility on the Level of Single-Electron Tunneling**, PRX 2012