



Entanglement on demand



Gershoni, Lindner, Akopian,
Berlitzky, Poem



On demand





$$|0\rangle \otimes |0\rangle + \varepsilon (|V\rangle \otimes |H\rangle + |H\rangle \otimes |V\rangle)$$

junk



Entangled piece

Quantum dots




$|\text{incoming photon}\rangle \rightarrow |\text{photon pair}\rangle$

Which path and interference




Monitoring the path kills interference

Which path and entanglement



Entanglement: A 2 photon analog of interference



Monitoring a cascade



Scales



classical correlations



Independence and correlations

independence

$$P_{ab}(x, y) = P_a(x)P_b(y)$$



Correlations due to common source

$$P_{ab}(x, y) = \sum_j p_j P_a^j(x) P_b^j(y)$$

Separable states

$$\rho_S = \sum p_j \rho_j^A \otimes \rho_j^B, \quad p_j > 0$$

Entangled states= Unseparable

$$\rho = |\psi\rangle\langle\psi|, \quad |\psi\rangle = \frac{|1\rangle \otimes |0\rangle - |0\rangle \otimes |1\rangle}{\sqrt{2}}$$

Negative probabilities

Peres test

$$\rho = \begin{pmatrix} A & B \\ B^* & C \end{pmatrix}, \quad \rho^P = \begin{pmatrix} A & B^* \\ B & C \end{pmatrix}$$

If transform has negative eigenvalue state is entangled

$$\rho = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad \rho^P = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}$$




Color as path monitor

Photons wave packet

$$|\psi\rangle = \alpha |HH\rangle \otimes \widehat{|p_H\rangle} + \beta |VV\rangle \otimes |p_V\rangle$$

$$\rho = \begin{pmatrix} |\alpha|^2 & 0 & 0 & \gamma \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \gamma & 0 & 0 & |\beta|^2 \end{pmatrix}$$

$$\gamma = \alpha\bar{\beta}\langle p_H|p_V\rangle$$




Monitors of the decay path kill the entanglement

Eliminating the monitor



$$|\psi\rangle \rightarrow (|HH\rangle + |VV\rangle) \otimes |p\rangle$$

$$\rho = \frac{1}{2} \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix}$$




Entanglement at a price

Fat is beautiful




Tomography


With monitor



Without monitor




Serendipity



ϵ_2

ϵ_1




The levels meander together with fixed splitting




Grad students Netanel Lindner and Nika Akopian
with Prof. Dudi Gershoni

Toshiba experiment

Forced degeneracy by annealing
And magnetic fields



Is degeneracy sufficient?



split


split

degenerate

Shields et. al. Nature 439 (2006)


Separable by Peres test

What went wrong



2 qubits Peres is iff

Octahedron=separable
Tetrahedron=all states
Cube= witnesses
Peres=reflection



Leinaas Myrheim Uvrom, Kenneth Avron

Local Operations

$$\dim \rho = 15$$

$$\rho \rightarrow (U_a \otimes U_b) \rho (U_a^\dagger \otimes U_b^\dagger)$$

$$U \in SU(2)$$

$$\dim(U_a \otimes U_b) = 9$$

$$U \in SL(2, \mathbb{C})$$

$$\dim(U_a \otimes U_b) = 12$$