

Deformations and Dilations of Chaotic Billiards: Dissipation Rate, and Quasiorthogonality of the Boundary Wave Functions

Alex Barnett, Doron Cohen, and Eric J. Heller

Department of Physics, Harvard University, Cambridge, Massachusetts 02138

(Received 9 March 2000)

We consider chaotic billiards in d dimensions, and study the matrix elements M_{nm} corresponding to general deformations of the boundary. We analyze the dependence of $|M_{nm}|^2$ on $\omega = (E_n - E_m)/\hbar$ using semiclassical considerations. This relates to an estimate of the energy dissipation rate when the deformation is periodic at frequency ω . We show that, for dilations and translations of the boundary, $|M_{nm}|^2$ vanishes like ω^4 as $\omega \rightarrow 0$, for rotations such as ω^2 , whereas for generic deformations it goes to a constant. Such special cases lead to quasiorthogonality of the eigenstates on the boundary.

PACS numbers: 05.45.Mt, 03.65.Sq

Chaotic cavities (billiards) in d dimensions are prototype systems for the study of classical chaos and its fingerprints on the properties of the quantum-mechanical eigenstates. As the properties of static billiards are beginning to be understood, questions naturally arise about deformations and their time dependence. It is perhaps not widely appreciated that certain deformations are very special, and that there is a close connection between the quantum and classical mechanics of such deformations in the case of ergodic systems. In this paper, which takes a fresh approach to these issues, we explore a special class of deformations which do not “heat” in the limit of small frequencies. We also establish a rather surprising relationship to a very successful numerical technique for finding billiard eigenfunctions.

We start with the one-particle Hamiltonian $\mathcal{H}_0(\mathbf{r}, \mathbf{p}) = \mathbf{p}^2/2m + V(\mathbf{r})$, where m is the particle mass, \mathbf{r} is the position of the particle inside the cavity, and \mathbf{p} is the conjugate momentum. We will take the limit $V(\mathbf{r}) \rightarrow \infty$ outside the cavity, zero otherwise, corresponding to Dirichlet boundary conditions. In this limit, the Hamiltonian is completely defined by the boundary shape. We call the volume of the cavity $\mathbf{V} \equiv \mathbf{L}^d$. Upon quantization a second length scale $\lambda_B \equiv 2\pi/k$ appears, where k is the wave number. For simple geometries the typical time between collisions with the walls is $\tau_{\text{col}} \sim L/v$, where v is the particle speed. The energy is $E = \frac{1}{2}mv^2$. Upon quantization the eigenenergies are $E_n = (\hbar k_n)^2/2m$.

A powerful tool for the classical analysis is known as the “Poincaré section.” Rather than following trajectories in the full (\mathbf{r}, \mathbf{p}) phase space, it is much more efficient to record only successive collisions with the boundary. This way we can deal with a canonical transformation (map) which is defined on a $2(d-1)$ -dimensional phase space. A similar idea is used in quantum mechanics: By using Green’s theorem, it is clear that all of the information about an eigenstate $\psi(\mathbf{r})$ is contained in the boundary normal derivative function $\varphi(\mathbf{s}) \equiv \mathbf{n} \cdot \nabla\psi$, where \mathbf{s} is a $(d-1)$ -dimensional coordinate on the boundary, and $\mathbf{n}(\mathbf{s})$ is the outward unit normal vector.

However, unlike the classical case, the reduction to the boundary is not satisfactory. One cannot define an associated Hilbert space that consists of the boundary functions. In particular, the orthogonality relation $\langle \psi_n | \psi_m \rangle = \delta_{nm}$ does not have an exact analog on the boundary. Still, the boundary functions “live” in an effective Hilbert space of dimension $\sim (L/\lambda_B)^{d-1}$, and it has been realized [1] that the following quasiorthogonality relation holds. Define an inner product

$$M_{nm} \equiv \frac{1}{2k^2} \oint \varphi_n(\mathbf{s}) \varphi_m(\mathbf{s}) (\mathbf{n} \cdot \hat{\mathbf{D}}) d\mathbf{s}, \quad (1)$$

where $\mathbf{D}(\mathbf{s}) = \mathbf{r}(\mathbf{s})$ is the displacement field corresponding to dilation (about an arbitrary origin), and $k_n \approx k_m \approx k$ [2]. It is well known that the normalization condition $\langle \psi_n | \psi_n \rangle = 1$ implies $M_{nn} = 1$. We give a proof of this exact result in the Appendix. On the other hand, the off-diagonal elements are only approximately zero [3].

The main purpose of this Letter is to study the band profile of the matrix M_{nm} for a general displacement field $\mathbf{D}(\mathbf{s})$. In particular we want to understand why, for special choices of $\mathbf{D}(\mathbf{s})$ (notably dilations), we have quasiorthogonality. Later we will explain that M_{nm} can be interpreted as the matrix element of a perturbation $\delta\mathcal{H}$ associated with a deformation of the boundary, such that $(\mathbf{n} \cdot \mathbf{D})\delta x$ is the normal displacement of a wall element, given a control parameter δx . In the following two paragraphs we explain the main motivations for our study.

The matrix elements $|M_{nm}|^2$ determine the rate of irreversible energy absorption by the particle (i.e., dissipation) due to external driving. Here “external driving” means time-dependent deformation of the boundary. Having exceptionally small $|M_{nm}|^2$ for special choices of $\mathbf{D}(\mathbf{s})$, such as dilations, translations, and rotations, implies an exceptionally small dissipation rate (“nonheating” effect). This observation goes against the naive kinetic picture that the rate of heating should not depend on how we “shake” the boundary. The special nature of translations and rotations for $\omega = 0$ has been recognized in the context of nuclear

dissipation [4,5]. Our present approach allows us to analyze the nonheating effect present for dilations as well, and provide the form of the low-frequency response of the system in all three cases (dilations, translations, and rotations).

There is another good motivation to study this issue. Recently, a powerful technique for finding clusters of billiard eigenstates and eigenenergies has been found by Vergini and Saraceno [1,6], with a speed typically $\sim 10^3$ greater than previous methods. This efficiency relies on the above quasiorthogonality relation, the associated numerical error being given by the deviation of M_{nm} from δ_{nm} . The authors tried to establish quasiorthogonality using the identity $M_{nm} = \delta_{nm} + [(k_m^2 - k_n^2)/2k^2]B_{nm}$, with $B_{nm} \equiv \langle \psi_n | \mathbf{r} \cdot \nabla | \psi_m \rangle$, and by assuming [7] that $|B_{nm}| \sim O(1)$. However, a naive random-wave argument would predict $|B_{nm}| \sim O(L/\lambda_B)^{(d-1)/2}$.

Figure 2 (below) displays the band profile $|M_{nm}|^2$ for three choices of the displacement field $\mathbf{D}(\mathbf{s})$. The band profile can be regarded as a function of either $\kappa = k_n - k_m$ or, equivalently, $\omega = (E_n - E_m)/\hbar$, related via $\omega = v\kappa$. The three band profiles differ in their peak structure, and also in their $\omega \rightarrow 0$ limits: notably for dilations $|M_{nm}|^2$ vanishes in this limit. Our aim is to understand the overall ω dependence, and the small ω behavior in particular. For the calculation of band profile we used all 451 eigenstates of the 2D quarter stadium (see Fig. 1) lying between $398 < k < 402$, found by using the method of Vergini and Saraceno [1]. For this particular chaotic shape a remarkably good basis set (size of order L/λ_B) of real and evanescent plane waves has been devised [6], which allows the *tension* error (defined as the boundary integral of ψ^2) to be typically 3×10^{-11} in our calculation, (maximum 2×10^{-10} for any state). The resulting errors in φ manifest themselves only when $|M_{nm}|^2$ reaches its lowest reliable value $\sim 10^{-10}$, visible as bottoming out in the leftmost point of the inset of Fig. 2.

In order to understand the quantum-mechanical band profile, we can first assume that the eigenstates look like uncorrelated random waves. A lengthy but straightforward calculation [8] leads to the result

$$|M_{nm}|^2 \approx \frac{2\langle |\cos(\theta)|^3 \rangle}{\Omega_d} \frac{\lambda_B^{d-1}}{V^2} \oint (\mathbf{n} \cdot \mathbf{D})^2 ds, \quad (2)$$

where the geometric factors for $d = 2, 3, \dots$ are $\Omega_d = 2\pi, 4\pi, \dots$ and $\langle |\cos(\theta)|^3 \rangle = 4/(3\pi), 1/4, \dots$. If the dis-




FIG. 1. The two-dimensional billiards under numerical study. Left: Bunimovich quarter stadium ($a/R = 1$). Right: generalized Sinai billiard ($a/b = 2$, $\theta_1 = 0.2$, $\theta_2 = 0.5$).

placement field is normalized such that $|\mathbf{D}| \sim L$, then we get $|M_{nm}|^2 \sim (\lambda_B/L)^{d-1}$. Note that the above result implies that $|M_{nm}|^2$ is independent of ω .

To go beyond the random-wave estimate (2), we adopt a more physically appealing point of view. We include a parametric deformation of the billiard shape via the Hamiltonian $\mathcal{H}(\mathbf{r}, \mathbf{p}; x) = \mathbf{p}^2/2m + V(\mathbf{r} - x\mathbf{D}(\mathbf{r}))$, where x controls the deformation. Note that the displacement field \mathbf{D} is regarded as a function of \mathbf{r} . The normal displacement of a wall element is $(\mathbf{n} \cdot \mathbf{D})x$. The position of a particle in the vicinity of a wall element is conveniently described by $Q = (\mathbf{s}, z)$, where \mathbf{s} is a surface coordinate and z is a perpendicular “radial” coordinate. We set $V(\mathbf{r}) = V_0$ outside the undeformed billiard; later we take the limit $V_0 \rightarrow \infty$. We have $\partial\mathcal{H}/\partial x = -[\mathbf{n}(\mathbf{s}) \cdot \mathbf{D}(\mathbf{s})]V_0\delta(z)$. The logarithmic derivative with respect to z of an eigenfunction on the boundary is $\varphi(\mathbf{s})/\psi(\mathbf{s})$. For $z > 0$ the wave function $\psi(\mathbf{r})$ is a decaying exponential. Hence the logarithmic derivative of the wave function on the boundary should be equal to $-\sqrt{2mV_0}/\hbar$. Consequently one obtains $(\partial\mathcal{H}/\partial x)_{nm} = -[(\hbar k)^2/m]M_{nm}$. Thus the band profile of M_{nm} is equal (up to a factor) to the band profile of the perturbation $\delta\mathcal{H}$ due to a deformation of the boundary (see also Refs. [5,8,9]).




FIG. 2. The band profile in the two-dimensional quarter stadium at $k \approx 400$ for three choices of deformation field: dilation (D), a generic deformation (G), and a generic deformation restricted to parallel displacement of the stadium upper edge (Gp). G and Gp are chosen to be volume preserving. In each case, the solid line is the average $|M_{nm}|^2$ (estimation error 10%) versus $\omega = v(k_n - k_m)$, with $v \mapsto 1$, and the dashed line is the semiclassical estimate [Eq. (4)] (estimation error 3%). We normalized G and Gp so that they share the same random-wave estimate [Eq. (2)] as D; this is shown as a horizontal dotted line. The system-specific peak due to bouncing-ball orbits is labeled (bb). The inset is a log-log plot with average $|M_{nm}|^2$ shown as points.

We can now use semiclassical considerations [10]. The application to the cavity example has been introduced in [8]. Here we summarize the recipe. One should generate a very long (ergodic) classical trajectory, and define for it the fluctuating quantity $\mathcal{F}(t) = -\partial\mathcal{H}(\mathbf{r}, \mathbf{p}; x)/\partial x|_{x=0}$, where the time dependence of \mathcal{F} is due to the trajectory $(\mathbf{r}(t), \mathbf{p}(t))$. Hence

$$\mathcal{F}(t) = \sum_{\text{col}} 2m\nu \cos(\theta_{\text{col}}) D_{\text{col}} \delta(t - t_{\text{col}}), \quad (3)$$

where t_{col} is the time of a collision, D_{col} stands for $\mathbf{n} \cdot \mathbf{D}$ at the point of the collision, and $\nu \cos(\theta_{\text{col}})$ is the normal component of the particle's collision velocity. If the deformation is volume preserving, then $\langle \mathcal{F}(t) \rangle = 0$, otherwise it is convenient to subtract the (constant) average value. Now one can calculate the correlation function $C(\tau)$ of the fluctuating quantity $\mathcal{F}(t)$, and its Fourier transform $\tilde{C}(\omega) \equiv \int C(\tau) \exp(i\omega\tau) d\tau$. The semiclassical estimate for the matrix element is

$$\left\langle \left| \left(\frac{\partial\mathcal{H}}{\partial x} \right)_{nm} \right|^2 \right\rangle \approx \frac{\Delta}{2\pi\hbar} \tilde{C}\left(\frac{E_n - E_m}{\hbar}\right), \quad (4)$$

where Δ is the mean level spacing. In practice, it is convenient, without loss of generality, to work with units such that in (3) the time t is measured in units of length, and we make the replacements $m \mapsto 1$ and $\nu \mapsto 1$. Then (4) can be cast into the form $\langle |M_{nm}|^2 \rangle \approx (\Delta_k/2\pi) \tilde{C}(\kappa)$, where Δ_k is the mean level spacing in k .

Figure 2 shows the excellent agreement between the actual band profile and that predicted by Eq. (4) for generic deformations and dilation. Note that there were no fitted parameters in this match. In all estimations of $\tilde{C}(\omega)$ we have used single trajectories of $\sim 10^6$ consecutive collisions.

Understanding the band profile of $|M_{nm}|^2$ has now been reduced to a matter of finding a classical theory for $\tilde{C}(\omega)$. If we assume that Eq. (3) is a train of uncorrelated impulses, then its power spectrum would be that of white noise, namely, $\tilde{C}(\omega) \approx \text{const}$. A straightforward calculation [8] then leads to the random-wave result (2) already presented. However, in reality there are correlations in this train, and therefore we should expect $\tilde{C}(\omega)$ to have some structure on the frequency scale $\omega \sim 1/\tau_{\text{col}}$. By looking at Fig. 2 we see that the white noise expectation is reasonably satisfied for one of the ‘‘generic’’ deformations (G), but not in the other two cases (D, R). We also see nonuniversal peaks at $\omega \sim 1/\tau_{\text{col}} \sim 1$. We now explain that for $\omega \ll 1/\tau_{\text{col}}$ there is total failure of the white noise result for dilations, as well as for translations and rotations, and discuss further complications that may arise if the billiard system is not strongly chaotic.

In Fig. 3 we display $\tilde{C}(\omega)$ for a different billiard shape, a generalized Sinai billiard (Fig. 1), chosen because it does not suffer from the nongeneric marginally stable orbits found in the quarter stadium. Here we see very convincing evidence that for small frequencies we have $\tilde{C}(\omega) \approx \text{const}$




FIG. 3. The classical power spectrum $\tilde{C}(\omega)$ for $\mathcal{F}(t)$ corresponding to a generic deformation (G), dilation (D), translation (T), and rotation (R), in the case of the generalized Sinai billiard with $m = \nu = 1$. The estimation error is 13% for G and R, and 20% for D and T. The two dotted lines show ω^2 and ω^4 frequency dependence, for purposes of comparison.

for generic deformation, $\tilde{C}(\omega) \propto \omega^4$ for dilation and translation, and $\tilde{C}(\omega) \propto \omega^2$ for rotation. Thus the white noise expectation is indeed satisfied in the $\omega \ll 1/\tau_{\text{col}}$ regime for generic deformations, but fails for dilations, translations, and rotations, for which $\tilde{C}(\omega) \rightarrow 0$ as $\omega \rightarrow 0$. This property is known (in the context of eigenvalue spectra) as ‘‘rigidity’’ [11]. It implies that the train of impulses is strongly correlated, a result which at first sight seems inconsistent with the assumption of chaotic motion. We will explain that there is no inconsistency here.

The quantity $\mathcal{F}(t) = -\partial\mathcal{H}/\partial x$ is related to $\dot{\mathbf{p}} = -\partial\mathcal{H}/\partial\mathbf{r} = -\nabla V$, the instantaneous force on the particle, by $\mathcal{F}(t) = -\mathbf{D}(\mathbf{r}) \cdot \dot{\mathbf{p}}$. For translations we have $\mathbf{D} = \mathbf{e}$, where \mathbf{e} is a constant vector that defines a direction in space. We can write $\mathcal{F}(t) = (d/dt)\mathcal{G}(t)$, where $\mathcal{G}(t) = -m\mathbf{e} \cdot \mathbf{r}$. A similar relation holds for dilation $\mathbf{D} = \mathbf{r}$ with $\mathcal{G}(t) = -\frac{1}{2}m\mathbf{r}^2$. It follows that $\tilde{C}(\omega) = \omega^4 \tilde{C}_G(\omega)$, where $\tilde{C}_G(\omega)$ is the power spectrum of $\mathcal{G}(t)$. Assuming that $\mathcal{G}(t)$, unlike $\mathcal{F}(t)$, is a generic fluctuating quantity that looks like white noise, it follows that $\tilde{C}(\omega)$ is generically characterized by ω^4 behavior for either translations or dilations. For rotations we have $\mathbf{D} = \mathbf{e} \times \mathbf{r}$, and we can write $\mathcal{F}(t) = (d/dt)\mathcal{G}(t)$, where $\mathcal{G}(t) = -\mathbf{e} \cdot (\mathbf{r} \times \mathbf{p})$ is a projection of the particle's angular momentum vector. Consequently $\tilde{C}(\omega) = \omega^2 \tilde{C}_G(\omega)$, and we expect $\tilde{C}(\omega)$ to be generically characterized by ω^2 behavior in the case of rotations.

In the previous paragraph we assumed that generic fluctuating quantities such as \mathbf{r}^2 and $\mathbf{e} \cdot \mathbf{r}$ and $\mathbf{e} \cdot (\mathbf{r} \times \mathbf{p})$, as well as $\mathcal{F}(t)$ for any generic deformations, have a white noise power spectrum as $\omega \rightarrow 0$. Obviously, this ‘‘white noise assumption’’ should be verified for any particular example. If the motion is *not* strongly chaotic, meaning that $C(\tau)$ decays like a power law (say, $1/\tau^{1-\gamma}$ with

$0 < \gamma < 1$) rather than an exponential, then the universal behavior is modified: we may have $\omega^{-\gamma}$ behavior for small frequencies. For a generic system, for instance, the generalized Sinai billiard, we do not have this complication. The stadium example on the other hand is nongeneric: the trajectory can remain in the marginally stable “bouncing-ball” orbit (between the top and bottom edges) for long times, with a probability scaling as a power law in time. Depending on the choice of $\mathbf{D}(\mathbf{r})$, this may manifest itself in $C(\tau)$. For example, in Fig. 2 the deformation Gp involves a parallel displacement of the upper edge, and the resulting sensitivity to the bouncing-ball orbit leads to large enhancement of the fluctuation intensity $\tilde{C}(\omega = 0)$, and is suggestive of singular $\omega^{-\gamma}$ behavior for small ω .

Finally, consider the time-dependent problem which is described by the Hamiltonian $\mathcal{H}(\mathbf{r}, \mathbf{p}; x(t))$. It is well known that under quite general circumstances the dissipation is Ohmic ($\propto \dot{x}^2$) (see [8,12], and references therein). If $x(t) = A \sin(\omega t)$, linear response theory gives the long-time heating rate $d\langle \mathcal{H} \rangle / dt = \mu \cdot \frac{1}{2}(\omega A)^2$. The dissipation coefficient μ is determined by the matrix elements of (4) (which up to a factor equals $|M_{nm}|^2$), and therefore μ is proportional to $\tilde{C}(\omega)$. Our results imply that μ vanishes in the limit $\omega \rightarrow 0$ for translations. One should not be surprised [4], since this follows from Galilean invariance: One can view the limit $\omega \rightarrow 0$ as corresponding to the special case of constant \dot{x} . For constant nonzero \dot{x} the particle(s) in the cavity accommodate their motion to the reference frame of the cavity, and there is no dissipation. A similar argument holds for rotations. On the other hand, it is somewhat surprising that the same conclusion holds for dilations (the only other shape-preserving deformation) as well. This observation, as far as we know, has not been introduced previously in the literature.

We thank Eduardo Vergini and Mike Haggerty for important discussions. This work was supported by ITAMP and the National Science Foundation.

Appendix.—There exist a couple of lengthy vector-identity proofs [9,13] of the normalization $M_{nn} = 1$ for the dilation case $\mathbf{D} = \mathbf{r}$, for $d = 2$. Here we present a physically illuminating alternative that works for arbitrary d . We use a phase-space-preserving definition of dilation operator $U(\alpha) \equiv \exp(i\alpha G/\hbar)$. It is generated by the Hermitian operator $G = \frac{1}{2}(\mathbf{r} \cdot \mathbf{p} + \mathbf{p} \cdot \mathbf{r})$. Applying this dilation to wave functions gives the expansion,

$$U(\alpha)\psi(\mathbf{r}) \approx \psi(\mathbf{r}) + \alpha[(d/2)\psi + \mathbf{r} \cdot \nabla\psi] + \mathcal{O}(\alpha^2). \quad (5)$$

The operator also has the effect $U^\dagger \mathbf{r} U = e^\alpha \mathbf{r}$ and $U^\dagger \mathbf{p} U = e^{-\alpha} \mathbf{p}$. Consider now any Hamiltonian $\mathcal{H}_0 = \mathbf{p}^2/(2m) + V(\mathbf{r})$. By defining the parameter-dependent

version $\mathcal{H}(\mathbf{r}, \mathbf{p}; \alpha) = U(\alpha)\mathcal{H}_0(\mathbf{r}, \mathbf{p})U(\alpha)^\dagger$, it is straightforward to obtain

$$\left. \frac{\partial \mathcal{H}}{\partial \alpha} \right|_{\alpha=0} = \frac{\mathbf{p}^2}{m} - \mathbf{r} \cdot \nabla V, \quad (6)$$

whose matrix elements in the case of the billiard potential are $(\partial \mathcal{H} / \partial \alpha)_{nm} = [(\hbar k)^2/m][\delta_{nm} - M_{nm}]$. Thus the nondiagonal terms are the same as those of the deformation $\mathbf{D} = \mathbf{r}$. The diagonal elements can be calculated directly by taking the limit $\alpha \rightarrow 0$ of the expression $(\langle U\psi | \mathcal{H}_0 | U\psi \rangle - \langle \psi | \mathcal{H}_0 | \psi \rangle) / \alpha$. By using (5) and the fact that $\langle \psi | \mathbf{r} \cdot \nabla | \psi \rangle = -d/2$, one can easily show that the result equals zero. From here it follows that $M_{nn} = 1$.

-
- [1] E. Vergini and M. Saraceno, Phys. Rev. E **52**, 2204 (1995).
 - [2] As is common in the “quantum chaos” literature, $k_n \approx k_m$ implies that $k_n - k_m$ is classically small, but still $n - m$ may be very large. For $n = m$, we define $k \equiv k_n$.
 - [3] For degenerate $k_n = k_m$, they are shown to be exactly zero in the dilation, translation, and rotation cases [9].
 - [4] J. Blocki, Y. Boneh, J. R. Nix, J. Randrup, M. Robel, A. J. Sierk, and W. J. Swiatecki, Ann. Phys. (N.Y.) **113**, 330 (1978).
 - [5] S. E. Koonin, R. L. Hatch, and J. Randrup, Nucl. Phys. **A283**, 87 (1977); S. E. Koonin and J. Randrup, Nucl. Phys. **A289**, 475 (1977).
 - [6] E. Vergini, Ph.D. thesis, Universidad de Buenos Aires, 1995.
 - [7] However, the behavior $B \sim |k_n - k_m|$ has since been observed numerically [E. Vergini (personal communication)], in agreement with our results.
 - [8] D. Cohen, Ann. Phys. (N.Y.) (to be published), cond-mat/9902168.
 - [9] M. V. Berry and M. Wilkinson, Proc. R. Soc. London A **392**, 15 (1984).
 - [10] M. Feingold and A. Peres, Phys. Rev. A **34**, 591 (1986); M. Feingold, D. Leitner, and M. Wilkinson, Phys. Rev. Lett. **66**, 986 (1991); M. Wilkinson, M. Feingold, and D. Leitner, J. Phys. A **24**, 175 (1991); M. Feingold, A. Gioletta, F. M. Izrailev, and L. Molinari, Phys. Rev. Lett. **70**, 2936 (1993).
 - [11] Namely, the sequence of times $\{t_{\text{col}}\}$ is analogous to the sequence of energies $\{\epsilon_n\}$ of chaotic spectra. Rigidity means that $\tilde{C}(\omega)$, analogous to Berry’s form factor $K(\tau)$, goes to zero at the origin. See M. V. Berry, Proc. R. Soc. London A **400**, 229 (1985).
 - [12] D. Cohen, Phys. Rev. Lett. **82**, 4951 (1999); in *Proceedings of the International School of Physics “Enrico Fermi,” Course CXLIII*, edited by G. Casati, I. Guarneri, and U. Smilansky (IOS Press, Amsterdam, 2000).
 - [13] P. A. Boasman, Nonlinearity **7**, 485 (1994).