

E451: Operating rotation matrices on spin states (2002A2)

Submitted by: Adam Reichenthal

The problem:

Find the state vector Ψ_m in the standard base for a spin polarized in the XY plane at angle $\varphi = 60^\circ$. Relate to the following cases:

- (1) Spin 1/2.
- (2) Spin 1 with circular polarization.
- (3) Spin 1 with linear polarization.

You must express the elements of the state vector in your final answer with $e^{i\pi/\text{integer}}$, $\sqrt{2}$ etc. In order to prevent losing all points because of an algebraic mistake, please write first the analytic formula and only after calculate it to receive your final answer.

The solution:

(1) We can receive any spin 1/2 polarization state by combining a rotation round the Y axis and a rotation round the Z axis. The result is:

$$|\vec{e}_{\theta,\varphi}\rangle = R(\varphi)R(\theta)|\uparrow\rangle = e^{-i\varphi S_z} e^{-i\theta S_y} |\uparrow\rangle \rightarrow \begin{pmatrix} e^{-i\varphi/2} \cos(\theta/2) \\ e^{i\varphi/2} \sin(\theta/2) \end{pmatrix}$$

The rotation action that has been done on the spin 1/2 up is a rotation by $\theta = \pi/2$ round the Y axis and after that a rotation by $\varphi = \pi/3$ round the Z axis. Therefore:

$$|\Psi_m\rangle = e^{-i(\pi/6)} \cos(\pi/4) |\uparrow\rangle + e^{i(\pi/6)} \sin(\pi/4) |\downarrow\rangle$$

Analytic result:

$$\Psi_m = \begin{pmatrix} e^{-i\frac{\pi}{6}} \cos(\frac{\pi}{4}) \\ e^{i\frac{\pi}{6}} \sin(\frac{\pi}{4}) \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} e^{-i\frac{\pi}{6}} \\ e^{i\frac{\pi}{6}} \end{pmatrix}$$

(2) We will use a similar technique as at (1) once again but this time using dim=3 rotation matrices to be able to take care of spin 1 and using the linear basis.

We can rotate the state $|\uparrow\rangle$ in order to get other circularly polarized states:

$$|\vec{e}_{\theta,\varphi}\rangle = U(\varphi\vec{e}_z)U(\theta\vec{e}_y)|\uparrow\rangle = e^{-i\varphi S_z} e^{-i\theta S_y} |\uparrow\rangle \rightarrow \begin{pmatrix} \frac{1}{2}(1 + \cos\theta)e^{-i\varphi} \\ \frac{1}{\sqrt{2}}\sin\theta \\ \frac{1}{2}(1 - \cos\theta)e^{i\varphi} \end{pmatrix}$$

The rotation action that has been done on the spin 1 up is a rotation by $\theta = \pi/2$ round the Y axis and after that a rotation by $\varphi = \pi/3$ round the Z axis. Therefore:

$$|\Psi_m\rangle = \frac{1}{2}(1 + \cos(\pi/2))e^{-i\pi/3} |\uparrow\rangle + \frac{1}{\sqrt{2}}\sin(\pi/2) |\updownarrow\rangle + \frac{1}{2}(1 - \cos(\pi/2))e^{i\pi/3} |\downarrow\rangle$$

Analytic result:

$$\Psi_m = \frac{1}{2} \begin{pmatrix} e^{-i\frac{\pi}{3}} \\ \sqrt{2} \\ e^{i\frac{\pi}{3}} \end{pmatrix}$$

(3) Using the same technique as in (2) this time with a rotation matrix that rotates the state $|\uparrow\rangle$ in order to get other linearly polarized states:

$$|\vec{e}_{\theta,\varphi}\rangle = U(\varphi\vec{e}_z)U(\theta\vec{e}_y)|\uparrow\rangle = e^{-i\varphi S_z} e^{-i\theta S_y} |\uparrow\rangle \rightarrow \begin{pmatrix} -\frac{1}{\sqrt{2}}\sin\theta e^{-i\varphi} \\ \cos\theta \\ \frac{1}{\sqrt{2}}\sin\theta e^{i\varphi} \end{pmatrix}$$

We receive:

$$|\Psi_m\rangle = -\frac{1}{\sqrt{2}}\sin(\pi/2)e^{-i\pi/3}|\uparrow\rangle + \cos(\pi/2)|\uparrow\rangle + \frac{1}{\sqrt{2}}\sin(\pi/2)e^{i\pi/3}|\downarrow\rangle$$

Analytic result:

$$\Psi_m = \frac{1}{\sqrt{2}} \begin{pmatrix} -e^{-i\frac{\pi}{3}} \\ 0 \\ e^{i\frac{\pi}{3}} \end{pmatrix}$$